arXiv:1003.3300v1 [math.NT] 17 Mar 2010

Identities of symmetry for generalized twisted Bernoulli polynomials twisted by ramified roots of unity

DAE SAN KIM

Abstract. We derive eight identities of symmetry in three variables related to generalized twisted Bernoulli polynomials and generalized twisted power sums, both of which are twisted by ramified roots of unity. All of these are new, since there have been results only about identities of symmetry in two variables. The derivations of identities are based on the p-adic integral expression of the generating function for the generalized twisted Bernoulli polynomials and the quotient of p-adic integrals that can be expressed as the exponential generating function for the generalized twisted power sums.

Key words : generalized twisted Bernoulli polynomial, generalized twisted power sum, Dirichlet character, ramified roots of unity, p-adic integral, identities of symmetry.

MSC2010:11B68;11S80;05A19.

1. Introduction and preliminaries

Let p be a fixed prime. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}_p$ will respectively denote the ring of p-adic integers, the field of p-adic rational numbers and the completion of the algebraic closure of $\mathbb{Q}_p$. Assume that $|\ |_p$ is the normalized absolute value of $\mathbb{C}_p$ such that $|p|_p = \frac{1}{p}$. The group Γ of all roots of unity of $\mathbb{C}_p$ is the direct product of its subgroups Γ_u(the subgroup of unramified roots of unity) and Γ_r(the subgroup of ramified roots of unity). Namely,

$$\Gamma = \Gamma_u \cdot \Gamma_r,\ \Gamma_u \cap \Gamma_r = \{1\},$$

where

$$\Gamma_u = \{\xi \in \mathbb{C}_p | \xi^r = 1 \ for\ some\ r \in \mathbb{Z}_{>0}\ with\ (r,p) = 1\},$$

$$\Gamma_r = \{\xi \in \mathbb{C}_p | \xi^{p^s} = 1 \ for\ some\ s \in \mathbb{Z}_{>0}\}.$$

Let d be a fixed positive integer. Then we let

$$X = X_d = \varprojlim_{N} \mathbb{Z}/dp^N\mathbb{Z},$$

and let $\pi : X \to \mathbb{Z}_p$ be the map given by the inverse limit of the natural maps

$$\mathbb{Z}/dp^N\mathbb{Z} \to \mathbb{Z}/p^N\mathbb{Z}.$$

If g is a function on $\mathbb{Z}_p$,we will use the same notation to denote the function $g \circ \pi$. Let $\chi : (\mathbb{Z}/d\mathbb{Z})^* \to \overline{\mathbb{Q}}^*$ be a (primitive) Dirichlet character of conductor d. Then it will be pulled back to X via the natural map $X \to \mathbb{Z}/d\mathbb{Z}$. Here we fix, once and for all, an imbedding $\overline{\mathbb{Q}} \to \mathbb{C}_p$, so that χ is regarded as a map of X to $\mathbb{C}_p$(cf.[8])

For a uniformly differentiable function $f : X \to \mathbb{C}_p$, the p-adic Volkenborn-type integral of f is defined(cf.[5]) by

$$\int_X f(z)d\mu(z) = \lim_{N\to\infty} \frac{1}{dP^N} \sum_{j=0}^{dp^N-1} f(j).$$

Then it is easy to see that

$$\int_X f(z+1)d\mu(z) = \int_X f(z)d\mu(z) + f'(0). \tag{1.1}$$

More generally, we deduce from (1.1) that, for any positive integer n,

$$\int_X f(z+n)d\mu(z) = \int_X f(z)d\mu(z) + \sum_{a=0}^{n-1} f'(a). \tag{1.2}$$

Throughout this paper, we let $\xi \in \Gamma_r$ be any fixed root of unity, and let

$$E = \{t \in \mathbb{C}_p||t|_p < p^{\frac{-1}{p-1}}\}. \tag{1.3}$$

Then, for each fixed $t \in E$, the function e^{zt} is analytic on $\mathbb{Z}_p$ and hence considered as a function on X and, by applying (1.2) to f with $f(z) = \chi(z)\xi^z e^{zt}$, we get the p-adic integral expression of the generating function for the generalized twisted Bernoulli numbers $B_{n,\chi,\xi}$ attached to χ and ξ;

$$\int_X \chi(z)\xi^z e^{zt}d\mu(z) = \frac{t}{\xi^d e^{dt}-1} \sum_{a=0}^{d-1} \chi(a)\xi^a e^{at} = \sum_{n=0}^{\infty} B_{n,\chi,\xi}\frac{t^n}{n!}(t \in E). \tag{1.4}$$

So we have the following p-adic integral expression of the generating function for the generalized twisted Bernoulli polynomials $B_{n,\chi.\xi}(x)$ attached to χ and ξ;

(1.5)
$$\int_X \chi(z)\xi^z e^{(x+z)t}d\mu(z) = \frac{te^{xt}}{\xi^d e^{dt}-1} \sum_{a=0}^{d-1} \chi(a)\xi^a e^{at} = \sum_{n=0}^{\infty} B_{n,\chi,\xi}(x)\frac{t^n}{n!}(t \in E, x \in \mathbb{Z}_p).$$

Also, from (1.1) we have:

$$\int_X \xi^z e^{zt}d\mu(z) = \frac{t}{\xi e^t - 1}. \tag{1.6}$$

Let $S_k(n;\chi,\xi)$ denote the kth generalized twisted power sum of the first $n+1$ nonnegative integers attached to χ and ξ, namely

$$S_k(n;\chi,\xi) = \sum_{a=0}^{n} \chi(a)\xi^a a^k = \chi(0)\xi^0 0^k + \chi(1)\xi^1 1^k + \cdots + \chi(n)\xi^n n^k. \tag{1.7}$$

From (1.4), (1.6), and (1.7), one easily derives the following identities: for $w \in \mathbb{Z}_{>0}$,

$$\frac{dw\int_X \chi(x)\xi^x e^{xt}d\mu(x)}{\int_X \xi^{dwy}e^{dwyt}d\mu(y)} = \frac{\xi^{dw}e^{dwt}-1}{\xi^d e^{dt}-1}\sum_{a=0}^{d-1}\chi(a)\xi^a e^{at} \tag{1.8}$$

$$= \sum_{a=0}^{dw-1}\chi(a)\xi^a e^{at} \tag{1.9}$$

$$= \sum_{k=0}^{\infty} S_k(dw-1;\chi,\xi)\frac{t^k}{k!}\ (t\in E). \tag{1.10}$$

In what follows, we will always assume that the p-adic integrals of the various (twisted) exponential functions on X are defined for $t \in E$(cf. (1.3)), and therefore it will not be mentioned.

[1], [2], [6], [9] and [10] are some of the previous works on identities of symmetry in two variables involving Bernoulli polynomials and power sums. For the brief history, one is referred to those papers. For the first time, the idea of [6] was extended in [4] to the case of three variables so as to yield many new identities with abundant symmetry. This added some new identities of symmetry even to the existing ones in two variables as well.

In this paper, we will produce 8 basic identities of symmetry in three variables w_1,w_2,w_3 related to generalized twisted Bernoulli polynomials and generalized twisted power sums, both of which are twisted by ramified roots of unity(i.e., p-power roots of unity)(cf. (4.8)-(4.11), (4.14)-(4.17). All of these seem to be new, since there have been results only about identities of symmetry in two variables in the literature ([7]). On the other hand, in [3] the measure introduced by Koblitz(cf.[8])was exploited in order to treat the unramified roots of unity case (i.e., the orders of the roots of unity are prime to p and the conductors of Dirichlet characters).

The following is stated as Theorem 4.5 and an example of the full six symmetries in the positive integers w_1,w_2,w_3.

$$w_1^{n-1}\sum_{k=0}^{n}\binom{n}{k}\sum_{a=0}^{dw_1-1}\chi(a)\xi^{aw_2w_3}B_{k,\chi,\xi^{w_1w_3}}(w_2y_1+\frac{w_2}{w_1}a)$$
$$\times S_{n-k}(dw_3-1;\chi,\xi^{w_1w_2})w_2^{n-k}w_3^{k-1}$$

$$= w_1^{n-1}\sum_{k=0}^{n}\binom{n}{k}\sum_{a=0}^{dw_1-1}\chi(a)\xi^{aw_2w_3}B_{k,\chi,\xi^{w_1w_2}}(w_3y_1+\frac{w_3}{w_1}a)$$
$$\times S_{n-k}(dw_2-1;\chi,\xi^{w_1w_3})w_3^{n-k}w_2^{k-1}$$

$$= w_2^{n-1}\sum_{k=0}^{n}\binom{n}{k}\sum_{a=0}^{dw_2-1}\chi(a)\xi^{aw_1w_3}B_{k,\chi,\xi^{w_2w_3}}(w_1y_1+\frac{w_1}{w_2}a)$$
$$\times S_{n-k}(dw_3-1;\chi,\xi^{w_1w_2})w_1^{n-k}w_3^{k-1}$$

$$= w_2^{n-1}\sum_{k=0}^{n}\binom{n}{k}\sum_{a=0}^{dw_2-1}\chi(a)\xi^{aw_1w_3}B_{k,\chi,\xi^{w_1w_2}}(w_3y_1+\frac{w_3}{w_2}a)$$
$$\times S_{n-k}(dw_1-1;\chi,\xi^{w_2w_3})w_3^{n-k}w_1^{k-1}$$

$$= w_3^{n-1}\sum_{k=0}^{n}\binom{n}{k}\sum_{a=0}^{dw_3-1}\chi(a)\xi^{aw_1w_2}B_{k,\chi,\xi^{w_2w_3}}(w_1y_1+\frac{w_1}{w_3}a)$$
$$\times S_{n-k}(dw_2-1;\chi,\xi^{w_1w_3})w_1^{n-k}w_2^{k-1}$$

$$= w_3^{n-1}\sum_{k=0}^{n}\binom{n}{k}\sum_{a=0}^{dw_3-1}\chi(a)\xi^{aw_1w_2}B_{k,\chi,\xi^{w_1w_3}}(w_2y_1+\frac{w_2}{w_3}a)$$
$$\times S_{n-k}(dw_1-1;\chi,\xi^{w_2w_3})w_2^{n-k}w_1^{k-1}.$$

The derivations of identities are based on the p-adic integral expression of the generalized twisted Bernoulli polynomials in (1.5) and the quotient of integrals in (1.8)-(1.10) that can be expressed as the exponential generating function for the generalized twisted power sums. These abundance of symmetries would not be unearthed if such p-adic integral representations had not been available. We indebted this idea to the paper [7].

2. SEVERAL TYPES OF QUOTIENTS OF p-ADIC INTEGRALS

Here we will introduce several types of quotients of p-adic integrals on X or X^3 from which some interesting identities follow owing to the built-in symmetries in w_1, w_2, w_3. In the following, w_1, w_2, w_3 are all positive integers and all of the explicit expressions of integrals in (2.2), (2.4), (2.6), and (2.8) are obtained from the identity in (1.4) and (1.6). To ease notations, from now on we suppress μ and denote, for example, $d\mu(x)$ and $d\mu(x_1)d\mu(x_2)d\mu(x_3)$ respectively simply by dx and dX.

(a) Type Λ_{23}^{i} (for $i = 0, 1, 2, 3$)
$I(\Lambda_{23}^{i})$
(2.1)

$$= \frac{d^i\int_{X^3}\chi(x_1)\chi(x_2)\chi(x_3)\xi^{w_2w_3x_1+w_1w_3x_2+w_1w_2x_3} \times e^{(w_2w_3x_1+w_1w_3x_2+w_1w_2x_3+w_1w_2w_3(\sum_{j=1}^{3-i}y_j))t}dX}{(\int_X \xi^{dw_1w_2w_3x_4}e^{dw_1w_2w_3x_4t}dx_4)^i}$$

(2.2)

$$=\frac{(w_1w_2w_3)^{2-i}t^{3-i}e^{w_1w_2w_3(\sum_{j=1}^{3-i}y_j)t}(\xi^{dw_1w_2w_3}e^{dw_1w_2w_3t}-1)^i}{(\xi^{dw_2w_3}e^{dw_2w_3t}-1)(\xi^{dw_1w_3}e^{dw_1w_3t}-1)(\xi^{dw_1w_2}e^{dw_1w_2t}-1)}$$
$$\times(\sum_{a=0}^{d-1}\chi(a)\xi^{aw_2w_3}e^{aw_2w_3t})(\sum_{a=0}^{d-1}\chi(a)\xi^{aw_1w_3}e^{aw_1w_3t})(\sum_{a=0}^{d-1}\chi(a)\xi^{aw_1w_2}e^{aw_1w_2t}).$$

(b) Type Λ_{13}^{i} (for $i = 0, 1, 2, 3$)

$I(\Lambda_{13}^{i})$

(2.3)

$$= \frac{d^{i}\int_{X^{3}}\chi(x_1)\chi(x_2)\chi(x_3)\xi^{w_1x_1+w_2x_2+w_3x_3}e^{(w_1x_1+w_2x_2+w_3x_3+w_1w_2w_3(\sum_{j=1}^{3-i}y_j))t}dX}{(\int_{X}\xi^{dw_1w_2w_3x_4}e^{dw_1w_2w_3x_4t}dx_4)^{i}}$$

(2.4)

$$= \frac{(w_1w_2w_3)^{1-i}t^{3-i}e^{w_1w_2w_3(\sum_{j=1}^{3-i}y_j)t}(\xi^{dw_1w_2w_3}e^{dw_1w_2w_3t}-1)^{i}}{(\xi^{dw_1}e^{dw_1t}-1)(\xi^{dw_2}e^{dw_2t}-1)(\xi^{dw_3e^{dw_3t}}-1)}$$

$$\times(\sum_{a=0}^{d-1}\chi(a)\xi^{aw_1}e^{aw_1t})(\sum_{a=0}^{d-1}\chi(a)\xi^{aw_2}e^{aw_2t})(\sum_{a=0}^{d-1}\chi(a)\xi^{aw_3}e^{aw_3t}).$$

(c-0) Type Λ_{12}^{0}
$I(\Lambda_{12}^{0})$

(2.5)

$$= \int_{X^{3}}\chi(x_1)\chi(x_2)\chi(x_3)\xi^{w_1x_1+w_2x_2+w_3x_3}e^{(w_1x_1+w_2x_2+w_3x_3+w_2w_3y+w_1w_3y+w_1w_2y)t}dX$$

(2.6)

$$= \frac{w_1w_2w_3t^{3}e^{(w_2w_3+w_1w_3+w_1w_2)yt}}{(\xi^{dw_1}e^{dw_1t}-1)(\xi^{dw_2}e^{dw_2t}-1)(\xi^{dw_3}e^{dw_3t}-1)}$$

$$\times(\sum_{a=0}^{d-1}\chi(a)\xi^{aw_1}e^{aw_1t})(\sum_{a=0}^{d-1}\chi(a)\xi^{aw_2}e^{aw_2t})(\sum_{a=0}^{d-1}\chi(a)\xi^{aw_3}e^{aw_3t}).$$

(c-1) Type Λ_{12}^{1}

$$I(\Lambda_{12}^{1}) = \frac{d^{3}\int_{X^{3}}\chi(x_1)\chi(x_2)\chi(x_3)\xi^{w_1x_1+w_2x_2+w_3x_3}e^{(w_1x_1+w_2x_2+w_3x_3)t}dX}{\int_{X^{3}}\xi^{d(w_2w_3z_1+w_1w_3z_2+w_1w_2z_3)}e^{d(w_2w_3z_1+w_1w_3z_2+w_1w_2z_3)t}dZ} \tag{2.7}$$

$$= \frac{(\xi^{dw_2w_3}e^{dw_2w_3t}-1)(\xi^{dw_1w_3}e^{dw_1w_3t}-1)(\xi^{dw_1w_2}e^{dw_1w_2t}-1)}{w_1w_2w_3(\xi^{dw_1}e^{dw_1t}-1)(\xi^{dw_2}e^{dw_2t}-1)(\xi^{dw_3}e^{dw_3t}-1)} \tag{2.8}$$

$$\times(\sum_{a=0}^{d-1}\chi(a)\xi^{aw_1}e^{aw_1t})(\sum_{a=0}^{d-1}\chi(a)\xi^{aw_2}e^{aw_2t})(\sum_{a=0}^{d-1}\chi(a)\xi^{aw_3}e^{aw_3t}).$$

All of the above p-adic integrals of various types are invariant under all permutations of w_1, w_2, w_3, as one can see either from p-adic integral representations in (2.1), (2.3), (2.5), and (2.7) or from their explicit evaluations in (2.2), (2.4), (2.6), and (2.8).

3. Identities for generalized twisted Bernoulli polynomials

All of the following results can be easily obtained from (1.5) and (1.8)-(1.10). First, let's consider Type Λ_{23}^{i}, for each $i = 0, 1, 2, 3$.

(a-0)

$$
\begin{aligned}
&I(\Lambda_{23}^0)\\
&=\int_X \chi(x_1)\xi^{w_2w_3x_1}e^{w_2w_3(x_1+w_1y_1)t}dx_1\int_X \chi(x_2)\xi^{w_1w_3x_2}e^{w_1w_3(x_2+w_2y_2)t}dx_2\\
&\qquad\qquad\times\int_X \chi(x_3)\xi^{w_1w_2x_3}e^{w_1w_2(x_3+w_3y_3)t}dx_3\\
&=(\sum_{k=0}^{\infty}\frac{B_{k,\chi,\xi^{w_2w_3}}(w_1y_1)}{k!}(w_2w_3t)^k)(\sum_{l=0}^{\infty}\frac{B_{l,\chi,\xi^{w_1w_3}}(w_2y_2)}{l!}(w_1w_3t)^l)\\
&\qquad\qquad\times(\sum_{m=0}^{\infty}\frac{B_{m,\chi,\xi^{w_1w_2}}(w_3y_3)}{m!}(w_1w_2t)^m)
\end{aligned}
$$

(3.1)

$$
\begin{aligned}
&=\sum_{n=0}^{\infty}(\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,\chi,\xi^{w_2w_3}}(w_1y_1)B_{l,\chi,\xi^{w_1w_3}}(w_2y_2)B_{m,\chi,\xi^{w_1w_2}}(w_3y_3)\\
&\qquad\qquad\times w_1^{l+m}w_2^{k+m}w_3^{k+l})\frac{t^n}{n!},
\end{aligned}
$$

where the inner sum is over all nonnegative integers k, l, m, with $k+l+m=n$, and

$$\binom{n}{k,l,m}=\frac{n!}{k!l!m!}.$$

(a-1) Here we write $I(\Lambda_{23}^1)$ in two different ways:

(1) $I(\Lambda_{23}^1)$

(3.2)

$$
\begin{aligned}
&=\frac{1}{w_3}\int_X \chi(x_1)\xi^{w_2w_3x_1}e^{w_2w_3(x_1+w_1y_1)t}dx_1\int_X \chi(x_2)\xi^{w_1w_3x_2}e^{w_1w_3(x_2+w_2y_2)t}dx_2\\
&\qquad\qquad\times\frac{dw_3\int_X \chi(x_3)\xi^{w_1w_2x_3}e^{w_1w_2x_3t}dx_3}{\int_X \xi^{dw_1w_2w_3x_4}e^{dw_1w_2w_3x_4t}dx_4}\\
&=\frac{1}{w_3}(\sum_{k=0}^{\infty}B_{k,\chi,\xi^{w_2w_3}}(w_1y_1)\frac{(w_2w_3t)^k}{k!})(\sum_{l=0}^{\infty}B_{l,\chi,\xi^{w_1w_3}}(w_2y_2)\frac{(w_1w_3t)^l}{l!})\\
&\qquad\qquad\times(\sum_{m=0}^{\infty}S_m(dw_3-1;\chi,\xi^{w_1w_2})\frac{(w_1w_2t)^m}{m!})
\end{aligned}
$$

(3.3)

$$
\begin{aligned}
&=\sum_{n=0}^{\infty}(\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,\chi,\xi^{w_2w_3}}(w_1y_1)B_{l,\chi,\xi^{w_1w_3}}(w_2y_2)\\
&\qquad\qquad\times S_m(dw_3-1;\chi,\xi^{w_1w_2})w_1^{l+m}w_2^{k+m}w_3^{k+l-1})\frac{t^n}{n!}.
\end{aligned}
$$

(2) Invoking (1.9), (3.2) can also be written as

$$
\begin{aligned}
&I(\Lambda_{23}^1)\\
&\quad= \frac{1}{w_3} \sum_{a=0}^{dw_3-1} \chi(a)\xi^{aw_1w_2} \int_X \chi(x_1)\xi^{w_2w_3x_1} e^{w_2w_3(x_1+w_1y_1)t} dx_1\\
&\qquad\qquad\qquad\times \int_X \chi(x_2)\xi^{w_1w_3x_2} e^{w_1w_3(x_2+w_2y_2+\frac{w_2}{w_3}a)t} dx_2\\
&\quad= \frac{1}{w_3} \sum_{a=0}^{dw_3-1} \chi(a)\xi^{aw_1w_2} (\sum_{k=0}^{\infty} B_{k,\chi,\xi^{w_2w_3}}(w_1y_1)\frac{(w_2w_3t)^k}{k!})\\
&\qquad\qquad\qquad\times (\sum_{l=0}^{\infty} B_{l,\chi,\xi^{w_1w_3}}(w_2y_2+\frac{w_2}{w_3}a)\frac{(w_1w_3t)^l}{l!})
\end{aligned}
$$

(3.4)

$$
\begin{aligned}
&= \sum_{n=0}^{\infty} (w_3^{n-1} \sum_{k=0}^{n} \binom{n}{k} B_{k,\chi,\xi^{w_2w_3}}(w_1y_1)\\
&\quad\times \sum_{a=0}^{dw_3-1} \chi(a)\xi^{aw_1w_2} B_{n-k,\chi,\xi^{w_1w_3}}(w_2y_2+\frac{w_2}{w_3}a) w_1^{n-k} w_2^k)\frac{t^n}{n!}.
\end{aligned}
$$

(a-2) Here we write $I(\Lambda_{23}^2)$ in three different ways:

(1) $I(\Lambda_{23}^2)$

(3.5)

$$
\begin{aligned}
&= \frac{1}{w_2w_3} \int_X \chi(x_1)\xi^{w_2w_3x_1} e^{w_2w_3(x_1+w_1y_1)t} dx_1 \times \frac{dw_2 \int_X \chi(x_2)\xi^{w_1w_3x_2} e^{w_1w_3x_2t} dx_2}{\int_X \xi^{dw_1w_2w_3x_4} e^{dw_1w_2w_3x_4t} dx_4}\\
&\qquad\qquad\qquad\times \frac{dw_3 \int_X \chi(x_3)\xi^{w_1w_2x_3} e^{w_1w_2x_3t} dx_3}{\int_X \xi^{dw_1w_2w_3x_4} e^{dw_1w_2w_3x_4t} dx_4}\\
&= \frac{1}{w_2w_3} (\sum_{k=0}^{\infty} B_{k,\chi,\xi^{w_2w_3}}(w_1y_1)\frac{(w_2w_3t)^k}{k!})(\sum_{l=0}^{\infty} S_l(dw_2-1;\chi,\xi^{w_1w_3})\frac{(w_1w_3t)^l}{l!})\\
&\qquad\qquad\qquad\times (\sum_{m=0}^{\infty} S_m(dw_3-1;\chi,\xi^{w_1w_2})\frac{(w_1w_2t)^m}{m!})
\end{aligned}
$$

(3.6)

$$
\begin{aligned}
&= \sum_{n=0}^{\infty} (\sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,\chi,\xi^{w_2w_3}}(w_1y_1) S_l(dw_2-1;\chi,\xi^{w_1w_3})\\
&\qquad\qquad\qquad\times S_m(dw_3-1;\chi,\xi^{w_1w_2}) w_1^{l+m} w_2^{k+m-1} w_3^{k+l-1})\frac{t^n}{n!}.
\end{aligned}
$$

(2) Invoking (1.9), (3.5) can also be written as

$I(\Lambda_{23}^2)$

$$
\begin{aligned}
&= \frac{1}{w_2w_3} \sum_{a=0}^{dw_2-1} \chi(a)\xi^{aw_1w_3} \int_X \chi(x_1)\xi^{w_2w_3x_1} e^{w_2w_3(x_1+w_1y_1+\frac{w_1}{w_2}a)t} dx_1\\
&\qquad\qquad\times \frac{dw_3 \int_X \chi(x_3)\xi^{w_1w_2x_3} e^{w_1w_2x_3t} dx_3}{\int_X \xi^{dw_1w_2w_3x_4} e^{dw_1w_2w_3x_4t} dx_4}
\end{aligned}
\tag{3.7}
$$

$$
\begin{aligned}
&= \frac{1}{w_2w_3}\sum_{a=0}^{dw_2-1}\chi(a)\xi^{aw_1w_3}\Big(\sum_{k=0}^{\infty}B_{k,\chi,\xi^{w_2w_3}}(w_1y_1+\frac{w_1}{w_2}a)\frac{(w_2w_3t)^k}{k!}\Big) \\
&\qquad \times\Big(\sum_{l=0}^{\infty}S_l(dw_3-1;\chi,\xi^{w_1w_2})\frac{(w_1w_2t)^l}{l!}\Big)
\end{aligned}
$$

$$
\begin{aligned}
&= \sum_{n=0}^{\infty}\Big(w_2^{n-1}\sum_{k=0}^{n}\binom{n}{k}\sum_{a=0}^{dw_2-1}\chi(a)\xi^{aw_1w_3}B_{k,\chi,\xi^{w_2w_3}}(w_1y_1+\frac{w_1}{w_2}a) \\
&\qquad \times S_{n-k}(dw_3-1;\chi,\xi^{w_1w_2})w_1^{n-k}w_3^{k-1}\Big)\frac{t^n}{n!}.
\end{aligned} \tag{3.8}
$$

(3) Invoking (1.9) once again, (3.7) can be written as

$$
\begin{aligned}
I(\Lambda_{23}^2) &= \frac{1}{w_2w_3}\sum_{a=0}^{dw_2-1}\chi(a)\xi^{aw_1w_3}\sum_{b=0}^{dw_3-1}\chi(b)\xi^{aw_1w_2} \\
&\qquad \times\int_X \chi(x_1)\xi^{w_2w_3x_1}e^{w_2w_3(x_1+w_1y_1+\frac{w_1}{w_2}a+\frac{w_1}{w_3}b)t}dx_1 \\
&= \frac{1}{w_2w_3}\sum_{a=0}^{dw_2-1}\chi(a)\xi^{aw_1w_3}\sum_{b=0}^{dw_3-1}\chi(b)\xi^{bw_1w_2} \\
&\qquad \times\Big(\sum_{n=0}^{\infty}B_{n,\chi,\xi^{w_2w_3}}(w_1y_1+\frac{w_1}{w_2}a+\frac{w_1}{w_3}b)\frac{(w_2w_3t)^n}{n!}\Big)
\end{aligned}
$$

$$
\begin{aligned}
&= \sum_{n=0}^{\infty}\Big((w_2w_3)^{n-1}\sum_{a=0}^{dw_2-1}\sum_{b=0}^{dw_3-1}\chi(ab)\xi^{w_1(aw_3+bw_2)} \\
&\qquad \times B_{n,\chi,\xi^{w_2w_3}}(w_1y_1+\frac{w_1}{w_2}a+\frac{w_1}{w_3}b)\Big)\frac{t^n}{n!}.
\end{aligned} \tag{3.9}
$$

(a-3)

$$
\begin{aligned}
I(\Lambda_{23}^3) &= \frac{1}{w_1w_2w_3}\times\frac{dw_1\int_X\chi(x_1)\xi^{w_2w_3x_1}e^{w_2w_3x_1t}dx_1}{\int_X\xi^{dw_1w_2w_3x_4}e^{dw_1w_2w_3x_4t}dx_4} \\
&\qquad \times\frac{dw_2\int_X\chi(x_2)\xi^{w_1w_3x_2}e^{w_1w_3x_2t}dx_2}{\int_X\xi^{dw_1w_2w_3x_4}e^{dw_1w_2w_3x_4t}dx_4}\times\frac{dw_3\int_X\chi(x_3)\xi^{w_1w_2x_3}e^{w_1w_2x_3t}dx_3}{\int_X\xi^{dw_1w_2w_3x_4}e^{dw_1w_2w_3x_4t}dx_4}
\end{aligned}
$$

$$
\begin{aligned}
&= \frac{1}{w_1w_2w_3}\Big(\sum_{k=0}^{\infty}S_k(dw_1-1;\chi,\xi^{w_2w_3})\frac{(w_2w_3t)^k}{k!}\Big)\Big(\sum_{l=0}^{\infty}S_l(dw_2-1;\chi,\xi^{w_1w_3})\frac{(w_1w_3t)^l}{l!}\Big) \\
&\qquad \times\Big(\sum_{m=0}^{\infty}S_m(dw_3-1;\chi,\xi^{w_1w_2})\frac{(w_1w_2t)^m}{m!}\Big)
\end{aligned}
$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty}(\sum_{k+l+m=n} \binom{n}{k,l,m} S_k(dw_1-1;\chi,\xi^{w_2w_3}) S_l(dw_2-1;\chi,\xi^{w_1w_3}) \\
&\qquad\qquad \times S_m(dw_3-1;\chi,\xi^{w_1w_2}) w_1^{l+m-1} w_2^{k+m-1} w_3^{k+l-1}) \frac{t^n}{n!}.
\end{aligned} \tag{3.10}$$

(b) For Type Λ_{13}^i $(i=0,1,2,3)$, we may consider the analogous things to the ones in (a-0), (a-1), (a-2), and (a-3). However, each of those can be obtained from the corresponding ones in (a-0), (a-1), (a-2), and (a-3). Indeed, if we substitute w_2w_3, w_1w_3, w_1w_2 respectively for w_1, w_2, w_3 in (2.1), this amounts to replacing t by $w_1w_2w_3t$ and ξ by $\xi^{w_1w_2w_3}$ in (2.3). So, upon replacing w_1, w_2, w_3 respectively by w_2w_3, w_1w_3, w_1w_2 , dividing by $(w_1w_2w_3)^n$, and replacing $\xi^{w_1w_2w_3}$ by ξ, in each of the expressions of (3.1), (3.3), (3.4), (3.6), (3.8)-(3.10), we will get the corresponding symmetric identities for Type Λ_{13}^i $(i=0,1,2,3)$.

(c-0)

$$\begin{aligned}
&I(\Lambda_{12}^0) \\
&= \int_X \chi(x_1)\xi^{w_1x_1} e^{w_1(x_1+w_2y)t} dx_1 \int_X \chi(x_2)\xi^{w_2x_2} e^{w_2(x_2+w_3y)t} dx_2 \\
&\qquad\qquad \times \int_X \chi(x_3)\xi^{w_3x_3} e^{w_3(x_3+w_1y)t} dx_3 \\
&= (\sum_{n=0}^{\infty} \frac{B_{k,\chi,\xi^{w_1}}(w_2y)}{k!}(w_1t)^k)(\sum_{l=0}^{\infty} \frac{B_{l,\chi,\xi^{w_2}}(w_3y)}{l!}(w_2t)^l)(\sum_{m=0}^{\infty} \frac{B_{m,\chi,\xi^{w_3}}(w_1y)}{m!}(w_3t)^m)
\end{aligned}$$

$$= \sum_{n=0}^{\infty}(\sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,\chi,\xi^{w_1}}(w_2y) B_{l,\chi,\xi^{w_2}}(w_3y) B_{m,\chi,\xi^{w_3}}(w_1y) w_1^k w_2^l w_3^m) \frac{t^n}{n!}. \tag{3.11}$$

(c-1)

$$\begin{aligned}
I(\Lambda_{12}^1) &= \frac{1}{w_1w_2w_3} \frac{dw_2 \int_X \chi(x_1)\xi^{w_1x_1} e^{w_1x_1t} dx_1}{\int_X \xi^{dw_1w_2z_3} e^{dw_1w_2z_3t} dz_3} \times \frac{dw_3 \int_X \chi(x_2)\xi^{w_2x_2} e^{w_2x_2t} dx_2}{\int_X \xi^{dw_2w_3z_1} e^{dw_2w_3z_1t} dz_1} \\
&\qquad \times \frac{dw_1 \int_X \chi(x_3)\xi^{w_3x_3} e^{w_3x_3t} dx_3}{\int_X \xi^{dw_3w_1z_2} e^{dw_3w_1z_2t} dz_2} \\
&= \frac{1}{w_1w_2w_3} (\sum_{k=0}^{\infty} S_k(dw_2-1;\chi,\xi^{w_1}) \frac{(w_1t)^k}{k!})(\sum_{l=0}^{\infty} S_l(dw_3-1;\chi,\xi^{w_2}) \frac{(w_2t)^l}{l!}) \\
&\qquad \times (\sum_{m=0}^{\infty} S_m(dw_1-1;\chi,\xi^{w_3}) \frac{(w_3t)^m}{m!})
\end{aligned}$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty}(\sum_{k+l+m=n} \binom{n}{k,l,m} S_k(dw_2-1;\chi,\xi^{w_1}) S_l(dw_3-1;\chi,\xi^{w_2}) \\
&\qquad\qquad \times S_m(dw_1-1;\chi,\xi^{w_3}) w_1^{k-1} w_2^{l-1} w_3^{m-1}) \frac{t^n}{n!}.
\end{aligned} \tag{3.12}$$

4. Main theorems

As we noted earlier in the last paragraph of Section 2, the various types of quotients of p-adic integrals are invariant under any permutation of w_1, w_2, w_3. So the corresponding expressions in Section 3 are also invariant under any permutation of w_1, w_2, w_3. Thus our results about identities of symmetry will be immediate consequences of this observation.

However, not all permutations of an expression in Section 3 yield distinct ones. In fact, as these expressions are obtained by permuting w_1, w_2, w_3 in a single one labeled by them, they can be viewed as a group in a natural manner and hence it is isomorphic to a quotient of S_3. In particular, the number of possible distinct expressions are $1, 2, 3$ or 6. (a-0), (a-1(1)), (a-1(2)), and (a-2(2)) give the full six identities of symmetry, (a-2(1)) and (a-2(3)) yield three identities of symmetry, and (c-0) and (c-1) give two identities of symmetry, while the expression in (a-3) yields no identities of symmetry.

Here we will just consider the cases of Theorems 4.4 and 4.8, leaving the others as easy exercises for the reader. As for the case of Theorem 4.4, in addition to (4.11)-(4.13), we get the following three ones:

$$\begin{aligned}\sum_{k+l+m=n}\binom{n}{k,l,m}&B_{k,\chi,\xi^{w_2w_3}}(w_1y_1)S_l(dw_3-1;\chi,\xi^{w_1w_2})\\&\times S_m(dw_2-1;\chi,\xi^{w_1w_3})w_1^{l+m}w_3^{k+m-1}w_2^{k+l-1},\end{aligned}\tag{4.1}$$

$$\begin{aligned}\sum_{k+l+m=n}\binom{n}{k,l,m}&B_{k,\chi,\xi^{w_1w_3}}(w_2y_1)S_l(dw_1-1;\chi,\xi^{w_2w_3})\\&\times S_m(dw_3-1;\chi,\xi^{w_1w_2})w_2^{l+m}w_1^{k+m-1}w_3^{k+l-1},\end{aligned}\tag{4.2}$$

$$\begin{aligned}\sum_{k+l+m=n}\binom{n}{k,l,m}&B_{k,\chi,\xi^{w_1w_2}}(w_3y_1)S_l(dw_2-1;\chi,\xi^{w_1w_3})\\&\times S_m(dw_1-1;\chi,\xi^{w_2w_3})w_3^{l+m}w_2^{k+m-1}w_1^{k+l-1}.\end{aligned}\tag{4.3}$$

But, by interchanging l and m, we see that (4.1), (4.2), and (4.3) are respectively equal to (4.11), (4.12), and (4.13).

As to Theorem 4.8, in addition to (4.17) and (4.18), we have:

$$\begin{aligned}\sum_{k+l+m=n}\binom{n}{k,l,m}&S_k(dw_2-1;\chi,\xi^{w_1})S_l(dw_3-1;\chi,\xi^{w_2})\\&\times S_m(dw_1-1;\chi,\xi^{w_3})w_1^{k-1}w_2^{l-1}w_3^{m-1},\end{aligned}\tag{4.4}$$

$$\begin{aligned}\sum_{k+l+m=n}\binom{n}{k,l,m}&S_k(dw_3-1;\chi,\xi^{w_2})S_l(dw_1-1;\chi,\xi^{w_3})\\&\times S_m(dw_2-1;\chi,\xi^{w_1})w_2^{k-1}w_3^{l-1}w_1^{m-1},\end{aligned}\tag{4.5}$$

$$\begin{aligned}\sum_{k+l+m=n}\binom{n}{k,l,m}&S_k(dw_3-1;\chi,\xi^{w_1})S_l(dw_2-1;\chi,\xi^{w_3})\\&\times S_m(dw_1-1;\chi,\xi^{w_2})w_1^{k-1}w_3^{l-1}w_2^{m-1},\end{aligned}\tag{4.6}$$

$$\begin{aligned}\sum_{k+l+m=n}\binom{n}{k,l,m}&S_k(dw_2-1;\chi,\xi^{w_3})S_l(dw_1-1;\chi,\xi^{w_2})\\&\times S_m(dw_3-1;\chi,\xi^{w_1})w_3^{k-1}w_2^{l-1}w_1^{m-1}.\end{aligned}\tag{4.7}$$

However, (4.4) and (4.5) are equal to (4.17), as we can see by applying the permutations $k \to l, l \to m, m \to k$ for (4.4) and $k \to m, l \to k, m \to l$ for (4.5). Similarly, we see that (4.6) and (4.7) are equal to (4.18), by applying permutations $k \to l, l \to m, m \to k$ for (4.6) and $k \to m, l \to k, m \to l$ for (4.7).

Theorem 4.1. *Let w_1, w_2, w_3 be any positive integers. Then we have:*

$$\begin{aligned}&\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,\chi,\xi^{w_2w_3}}(w_1y_1)B_{l,\chi,\xi^{w_1w_3}}(w_2y_2)\\&\qquad\times B_{m,\chi,\xi^{w_1w_2}}(w_3y_3)w_1^{l+m}w_2^{k+m}w_3^{k+l}\\
&=\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,\chi,\xi^{w_2w_3}}(w_1y_1)B_{l,\chi,\xi^{w_1w_2}}(w_3y_2)\\&\qquad\times B_{m,\chi,\xi^{w_1w_3}}(w_2y_3)w_1^{l+m}w_3^{k+m}w_2^{k+l}\\
&=\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,\chi,\xi^{w_1w_3}}(w_2y_1)B_{l,\chi,\xi^{w_2w_3}}(w_1y_2)\\&\qquad\times B_{m,\chi,\xi^{w_1w_2}}(w_3y_3)w_2^{l+m}w_1^{k+m}w_3^{k+l}\\
&=\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,\chi,\xi^{w_1w_3}}(w_2y_1)B_{l,\chi,\xi^{w_1w_2}}(w_3y_2)\\&\qquad\times B_{m,\chi,\xi^{w_2w_3}}(w_1y_3)w_2^{l+m}w_3^{k+m}w_1^{k+l}\\
&=\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,\chi,\xi^{w_1w_2}}(w_3y_1)B_{l,\chi,\xi^{w_2w_3}}(w_1y_2)\\&\qquad\times B_{m,\chi,\xi^{w_1w_3}}(w_2y_3)w_3^{l+m}w_1^{k+m}w_2^{k+l}\\
&=\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,\chi,\xi^{w_1w_2}}(w_3y_1)B_{l,\chi,\xi^{w_1w_3}}(w_2y_2)\\&\qquad\times B_{m,\chi,\xi^{w_2w_3}}(w_1y_3)w_3^{l+m}w_2^{k+m}w_1^{k+l}.\end{aligned}\tag{4.8}$$

Theorem 4.2. *Let w_1, w_2, w_3 be any positive integers. Then we have:*

$$\begin{aligned}&\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,\chi,\xi^{w_2w_3}}(w_1y_1)B_{l,\chi,\xi^{w_1w_3}}(w_2y_2)\\&\qquad\times S_m(dw_3-1;\chi,\xi^{w_1w_2})w_1^{l+m}w_2^{k+m}w_3^{k+l-1}\\
&=\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,\chi,\xi^{w_2w_3}}(w_1y_1)B_{l,\chi,\xi^{w_1w_2}}(w_3y_2)\\&\qquad\times S_m(dw_2-1;\chi,\xi^{w_1w_3})w_1^{l+m}w_3^{k+m}w_2^{k+l-1}\end{aligned}$$

$$\begin{aligned} &= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,\chi,\xi^{w_1w_3}}(w_2y_1) B_{l,\chi,\xi^{w_2w_3}}(w_1y_2) \\ &\qquad \times S_m(dw_3-1;\chi,\xi^{w_1w_2}) w_2^{l+m} w_1^{k+m} w_3^{k+l-1} \\ &= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,\chi,\xi^{w_1w_3}}(w_2y_1) B_{l,\chi,\xi^{w_1w_2}}(w_3y_2) \\ &\qquad \times S_m(dw_1-1;\chi,\xi^{w_2w_3}) w_2^{l+m} w_3^{k+m} w_1^{k+l-1} \\ &= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,\chi,\xi^{w_1w_2}}(w_3y_1) B_{l,\chi,\xi^{w_1w_3}}(w_2y_2) \\ &\qquad \times S_m(dw_1-1;\chi,\xi^{w_2w_3}) w_3^{l+m} w_2^{k+m} w_1^{k+l-1} \\ &= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,\chi,\xi^{w_1w_2}}(w_3y_1) B_{l,\chi,\xi^{w_2w_3}}(w_1y_2) \\ &\qquad \times S_m(dw_2-1;\chi,\xi^{w_1w_3}) w_3^{l+m} w_1^{k+m} w_2^{k+l-1}. \end{aligned} \tag{4.9}$$

Theorem 4.3. *Let w_1, w_2, w_3 be any positive integers. Then we have:*

$$\begin{aligned} & w_1^{n-1} \sum_{k=0}^{n} \binom{n}{k} B_{k,\chi,\xi^{w_1w_2}}(w_3y_1) \sum_{a=0}^{dw_1-1} \chi(a)\xi^{aw_2w_3} B_{n-k,\chi,\xi^{w_1w_3}}(w_2y_2 + \frac{w_2}{w_1}a) w_3^{n-k} w_2^{k} \\ =& w_1^{n-1} \sum_{k=0}^{n} \binom{n}{k} B_{k,\chi,\xi^{w_1w_3}}(w_2y_1) \sum_{a=0}^{dw_1-1} \chi(a)\xi^{aw_2w_3} B_{n-k,\chi,\xi^{w_1w_2}}(w_3y_2 + \frac{w_3}{w_1}a) w_2^{n-k} w_3^{k} \\ =& w_2^{n-1} \sum_{k=0}^{n} \binom{n}{k} B_{k,\chi,\xi^{w_1w_2}}(w_3y_1) \sum_{a=0}^{dw_2-1} \chi(a)\xi^{aw_1w_3} B_{n-k,\chi,\xi^{w_2w_3}}(w_1y_2 + \frac{w_1}{w_2}a) w_3^{n-k} w_1^{k} \\ =& w_2^{n-1} \sum_{k=0}^{n} \binom{n}{k} B_{k,\chi,\xi^{w_2w_3}}(w_1y_1) \sum_{a=0}^{dw_2-1} \chi(a)\xi^{aw_1w_3} B_{n-k,\chi,\xi^{w_1w_2}}(w_3y_2 + \frac{w_3}{w_2}a) w_1^{n-k} w_3^{k} \\ =& w_3^{n-1} \sum_{k=0}^{n} \binom{n}{k} B_{k,\chi,\xi^{w_1w_3}}(w_2y_1) \sum_{a=0}^{dw_3-1} \chi(a)\xi^{aw_1w_2} B_{n-k,\chi,\xi^{w_2w_3}}(w_1y_2 + \frac{w_1}{w_2}a) w_2^{n-k} w_1^{k} \\ =& w_3^{n-1} \sum_{k=0}^{n} \binom{n}{k} B_{k,\chi,\xi^{w_2w_3}}(w_1y_1) \sum_{a=0}^{dw_3-1} \chi(a)\xi^{aw_1w_2} B_{n-k,\chi,\xi^{w_1w_3}}(w_2y_2 + \frac{w_2}{w_3}a) w_1^{n-k} w_2^{k}. \end{aligned} \tag{4.10}$$

Theorem 4.4. *Let w_1, w_2, w_3 be any positive integers. Then we have the following three symmetries in w_1, w_2, w_3:*

$$\begin{aligned} & \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,\chi,\xi^{w_2w_3}}(w_1y_1) S_l(dw_2-1;\chi,\xi^{w_1w_3}) \\ &\qquad \times S_m(dw_3-1;\chi,\xi^{w_1w_2}) w_1^{l+m} w_2^{k+m-1} w_3^{k+l-1} \end{aligned} \tag{4.11}$$

$$\begin{aligned} &= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,\chi,\xi^{w_1w_3}}(w_2y_1) S_l(dw_3-1;\chi,\xi^{w_1w_2}) \\ &\qquad \times S_m(dw_1-1;\chi,\xi^{w_2w_3}) w_2^{l+m} w_3^{k+m-1} w_1^{k+l-1} \end{aligned} \tag{4.12}$$

$$\begin{aligned}&= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,\chi,\xi^{w_1w_2}}(w_3y_1) S_l(dw_1-1,\chi,\xi^{w_2w_3}) \\ &\qquad \times S_m(dw_2-1;\chi,\xi^{w_1w_3}) w_3^{l+m} w_1^{k+m-1} w_2^{k+l-1}.\end{aligned} \tag{4.13}$$

Theorem 4.5. *Let w_1, w_2, w_3 be any positive integers. Then we have:*

$$\begin{aligned}& w_1^{n-1} \sum_{k=0}^{n} \binom{n}{k} \sum_{a=0}^{dw_1-1} \chi(a)\xi^{aw_2w_3} B_{k,\chi,\xi^{w_1w_3}}(w_2y_1 + \frac{w_2}{w_1}a) \\ &\qquad \times S_{n-k}(dw_3-1;\chi,\xi^{w_1w_2}) w_2^{n-k} w_3^{k-1}\end{aligned} \tag{4.14}$$

$$\begin{aligned}&= w_1^{n-1} \sum_{k=0}^{n} \binom{n}{k} \sum_{a=0}^{dw_1-1} \chi(a)\xi^{aw_2w_3} B_{k,\chi,\xi^{w_1w_2}}(w_3y_1 + \frac{w_3}{w_1}a) \\ &\qquad \times S_{n-k}(dw_2-1;\chi,\xi^{w_1w_3}) w_3^{n-k} w_2^{k-1} \\ &= w_2^{n-1} \sum_{k=0}^{n} \binom{n}{k} \sum_{a=0}^{dw_2-1} \chi(a)\xi^{aw_1w_3} B_{k,\chi,\xi^{w_2w_3}}(w_1y_1 + \frac{w_1}{w_2}a) \\ &\qquad \times S_{n-k}(dw_3-1;\chi,\xi^{w_1w_2}) w_1^{n-k} w_3^{k-1} \\ &= w_2^{n-1} \sum_{k=0}^{n} \binom{n}{k} \sum_{a=0}^{dw_2-1} \chi(a)\xi^{aw_1w_3} B_{k,\chi,\xi^{w_1w_2}}(w_3y_1 + \frac{w_3}{w_2}a) \\ &\qquad \times S_{n-k}(dw_1-1;\chi,\xi^{w_2w_3}) w_3^{n-k} w_1^{k-1} \\ &= w_3^{n-1} \sum_{k=0}^{n} \binom{n}{k} \sum_{a=0}^{dw_3-1} \chi(a)\xi^{aw_1w_2} B_{k,\chi,\xi^{w_2w_3}}(w_1y_1 + \frac{w_1}{w_3}a) \\ &\qquad \times S_{n-k}(dw_2-1;\chi,\xi^{w_1w_3}) w_1^{n-k} w_2^{k-1} \\ &= w_3^{n-1} \sum_{k=0}^{n} \binom{n}{k} \sum_{a=0}^{dw_3-1} \chi(a)\xi^{aw_1w_2} B_{k,\chi,\xi^{w_1w_3}}(w_2y_1 + \frac{w_2}{w_3}a) \\ &\qquad \times S_{n-k}(dw_1-1;\chi,\xi^{w_2w_3}) w_2^{n-k} w_1^{k-1}.\end{aligned}$$

Theorem 4.6. *Let w_1, w_2, w_3 be any positive integers. Then we have the following three symmetries in w_1, w_2, w_3:*

$$\begin{aligned}&(w_1w_2)^{n-1} \sum_{a=0}^{dw_1-1} \sum_{b=0}^{dw_2-1} \chi(ab)\xi^{w_3(aw_2+bw_1)} B_{n,\chi,\xi^{w_1w_2}}(w_3y_1 + \frac{w_3}{w_1}a + \frac{w_3}{w_2}b) \\ =&(w_2w_3)^{n-1} \sum_{a=0}^{dw_2-1} \sum_{b=0}^{dw_3-1} \chi(ab)\xi^{w_1(aw_3+bw_2)} B_{n,\chi,\xi^{w_2w_3}}(w_1y_1 + \frac{w_1}{w_2}a + \frac{w_1}{w_3}b) \\ =&(w_3w_1)^{n-1} \sum_{a=0}^{dw_3-1} \sum_{b=0}^{dw_1-1} \chi(ab)\xi^{w_2(aw_1+bw_3)} B_{n,\chi,\xi^{w_1w_3}}(w_2y_1 + \frac{w_2}{w_3}a + \frac{w_2}{w_1}b).\end{aligned} \tag{4.15}$$

Theorem 4.7. *Let w_1, w_2, w_3 be any positive integers. Then we have the following two symmetries in w_1, w_2, w_3:*

$$\begin{aligned}&\sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,\chi,\xi^{w_3}}(w_1 y) B_{l,\chi,\xi^{w_1}}(w_2 y) B_{m,\chi,\xi^{w_2}}(w_3 y) w_3^k w_1^l w_2^m \\ &= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,\chi,\xi^{w_2}}(w_1 y) B_{l,\chi,\xi^{w_1}}(w_3 y) B_{m,\chi,\xi^{w_3}}(w_2 y) w_2^k w_1^l w_3^m. \end{aligned} \tag{4.16}$$

Theorem 4.8. *Let w_1, w_2, w_3 be any positive integers. Then we have the following two symmetries in w_1, w_2, w_3:*

$$\begin{aligned}&\sum_{k+l+m=n} \binom{n}{k,l,m} S_k(dw_1 - 1; \chi, \xi^{w_3}) S_l(dw_2 - 1; \chi, \xi^{w_1}) \\ &\qquad \times S_m(dw_3 - 1; \chi, \xi^{w_2}) w_3^{k-1} w_1^{l-1} w_2^{m-1} \end{aligned} \tag{4.17}$$

$$\begin{aligned}&= \sum_{k+l+m=n} \binom{n}{k,l,m} S_k(dw_1 - 1; \chi, \xi^{w_2}) S_l(dw_3 - 1; \chi, \xi^{w_1}) \\ &\qquad \times S_m(dw_2 - 1; \chi, \xi^{w_3}) w_2^{k-1} w_1^{l-1} w_3^{m-1}. \end{aligned} \tag{4.18}$$

References

1. E. Deeba and D. Rodriguez, *Stirling's and Bernoulli numbers* , Amer. Math. Monthly . **98** (1991), 423-426.
2. F. T. Howard, *Applications of a recurrence for the Bernoulli numbers,* J. Number Theory **52**(1995), 157-172.
3. D. S. Kim, *Identities of symmetry for generalized twisted Bernoulli polynomials twisted by unramified roots of unity,* submitted.
4. D. S. Kim and K. H. Park, *Identities of symmetry for Bernoulli polynomials arising from quotients of Volkenborn integrals invariant under S_3*, submitted.
5. T. Kim, *On a q-analogue of the p-adic log gamma functions and related integrals,* J. Number Theory **76** (1999), 320-329.
6. T. Kim, *Symmetry p-adic invariant integral on $\mathbb{Z}_p$ for Bernoulli and Euler polynomials,* J. Difference Equ. Appl. **14** (2008), 1267-1277.
7. T. Kim, *On the symmetric properties for the generalized twisted Bernoulli polynomials,* J. Inequal. Appl. 2009(2009), 8 pages(Article ID 164743).
8. N. Koblitz, *A new proof of certain formulas for p-adic L-functions,* Duke Math. J. **46** (1979), 455-468.
9. H. Tuenter, *A symmetry of power sum polynomials and Bernoulli numbers,* Amer. Math. Monthly **108** (2001), 258-261.
10. S. Yang, *An identity of symmetry for the Bernoulli polynomials,* Discrete Math. **308** (2008), 550-554.

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 121-742, KOREA
E-mail address: `dskim@sogong.ac.kr`